

Convergent variational calculation of positronium-hydrogen-atom scattering lengths

Sadhan K Adhikari[†] and Puspajit Mandal^{†,§}

[†]*Instituto de Física Teórica, Universidade Estadual Paulista 01.405-900 São Paulo, São Paulo, Brazil*

[§]*Department of Mathematics, Visva-Bharati, Santiniketan 731 235, India*

(February 2, 2008)

Abstract

We present a convergent variational basis-set calculational scheme for elastic scattering of positronium atom by hydrogen atom in S wave. Highly correlated trial functions with appropriate symmetry are needed for achieving convergence. We report convergent results for scattering lengths in atomic units for both singlet ($= 3.49 \pm 0.20$) and triplet ($= 2.46 \pm 0.10$) states.

PACS Number(s): 34.90.+q, 36.10.Dr

Lately, there has been interest in the experimental [1] and theoretical [2–8] studies of ortho positronium (Ps) atom scattering by different neutral atomic and molecular targets. The Ps-H system is theoretically the most simple and fundamental and a complete understanding of this system is necessary before a venture to more complex targets [7–9]. There have been R-matrix [3], close-coupling (CC) [4,10] and model-potential [5,11] calculations for Ps-H scattering. Here we present a convergent variational basis-set calculational scheme for low-energy Ps-H scattering in S wave below the lowest Ps-excitation threshold at 5.1 eV. Using this method we report numerical results for scattering length of electronic singlet and triplet states.

A recent study based on a regularized nonlocal electron-exchange model potential [6] yielded low-energy (total) cross sections in agreement with experiment for Ps scattering by He [6], Ne [7], Ar [7] and H₂ [8]. For the Ps-H system the model-potential results for S-wave singlet binding and resonance energies are in agreement with accurate variational estimates [12]. It would be interesting to see if the model-potential result for the Ps-H singlet scattering length agrees with the present convergent calculation.

Because of the existence of two identical fermions (electrons) in the Ps-H system, one needs to antisymmetrize the full wave function before attempting a solution of the scattering problem. The position vectors of the electrons – $\mathbf{r}_1$ (Ps) and $\mathbf{r}_2$ (H) – and positron ($\mathbf{x}$) with respect to (w.r.t.) the massive proton at the origin are as shown in figure 1. We also use the position vectors $\mathbf{s}_j = (\mathbf{x} + \mathbf{r}_j)/2$, $\rho_j = \mathbf{x} - \mathbf{r}_j$, $j = 1, 2$, $\mathbf{r}_{12} = \mathbf{r}_1 - \mathbf{r}_2$. The fully antisymmetric state $\psi_{\mathbf{k}}^A$ of Ps-H scattering is given by $|\psi_{\mathbf{k}}^A\rangle = \mathcal{A}_1|\psi_{\mathbf{k}}^1\rangle = (1 \pm P_{12})|\psi_{\mathbf{k}}^1\rangle = |\psi_{\mathbf{k}}^1\rangle \pm |\psi_{\mathbf{k}}^2\rangle$ where $\mathbf{k}$ is the incident momentum, the antisymmetrizer $\mathcal{A}_1$ is $(1 + P_{12})$ for the singlet state and $(1 - P_{12})$ for the triplet state with P_{12} the permutation operator of electrons 1 and 2. The function $\psi_{\mathbf{k}}^1$ refers to the Ps-H wave function with electron 1 forming the Ps as in figure 1 and $\psi_{\mathbf{k}}^2$ refers to the same with the two electrons interchanged.

The full Ps-H Hamiltonian H can be broken in two convenient forms as follows $H = H_1 + V_1 = H_2 + V_2$ where H_1 includes the full kinetic energy and intracuster interaction of H and Ps for the arrangement shown in figure 1 and V_1 is the sum of the intercluster interaction between H and Ps in the same configuration, H_2 and V_2 refer to the same quantities with the two electrons interchanged:

$$V_1 = \left[\frac{1}{x} - \frac{1}{r_1} + \frac{1}{r_{12}} - \frac{1}{\rho_2} \right], \quad V_2 = P_{12}V_1 = \left[\frac{1}{x} - \frac{1}{r_2} + \frac{1}{r_{12}} - \frac{1}{\rho_1} \right]. \quad (1)$$

The fully antisymmetric state satisfies the Lippmann-Schwinger equation [13]

$$|\psi_{\mathbf{k}}^A\rangle = |\phi_{\mathbf{k}}^1\rangle + G_1 V_1 |\psi_{\mathbf{k}}^A\rangle. \quad (2)$$

where the channel Green's function $G_1 \equiv (E + i0 - H_1)^{-1}$ and the incident wave $|\phi_{\mathbf{k}}^1\rangle$ satisfies $(E - H_1)|\phi_{\mathbf{k}}^1\rangle = 0$. The incident Ps energy $E = 6.8k^2$ eV. We are using atomic units (au) in which $a_0 = e = m = \hbar = 1$, where e (m) is the electronic charge (mass) and a_0 the Bohr radius. Using the definition of the antisymmetrized state we rewrite (2) as [13]

$$|\psi_{\mathbf{k}}^1\rangle = |\phi_{\mathbf{k}}^1\rangle + G_1 M_1 |\psi_{\mathbf{k}}^1\rangle \quad (3)$$

$$M_1 = V_1 \mathcal{A}_1 + (E - H_1)(1 - \mathcal{A}_1) \equiv \mathcal{A}_1 V_1 + (1 - \mathcal{A}_1)(E - H_1). \quad (4)$$

The properly symmetrized transition matrix for elastic scattering is defined by $\langle \phi_{\mathbf{k}}^1 | T^A | \phi_{\mathbf{k}}^1 \rangle = \langle \phi_{\mathbf{k}}^1 | V_1 | \psi_{\mathbf{k}}^A \rangle = \langle \phi_{\mathbf{k}}^1 | V_1 \mathcal{A}_1 | \psi_{\mathbf{k}}^1 \rangle = \langle \psi_{\mathbf{k}}^1 | \mathcal{A}_1 V_1 | \phi_{\mathbf{k}}^1 \rangle$ [13]. A basis-set calculational scheme for the transition matrix can be obtained from the following expression [14]

$$\langle \phi_{\mathbf{k}}^1 | T^A | \phi_{\mathbf{k}}^1 \rangle = \langle \psi_{\mathbf{k}}^1 | \mathcal{A}_1 V_1 | \phi_{\mathbf{k}}^1 \rangle + \langle \phi_{\mathbf{k}}^1 | \mathcal{A}_1 V_1 | \psi_{\mathbf{k}}^1 \rangle - \langle \psi_{\mathbf{k}}^1 | \mathcal{A}_1 V_1 - M_1 G_1 \mathcal{A}_1 V_1 | \psi_{\mathbf{k}}^1 \rangle. \quad (5)$$

Using (3), it can be verified that (5) is an identity if exact scattering wave functions $\psi_{\mathbf{k}}^1$ are used. If approximate wave functions are used, (5) is stationary w.r.t. small variations of $|\psi_{\mathbf{k}}^1\rangle$ but not with $\langle \psi_{\mathbf{k}}^1|$. This one-sided variational property emerges because of the lack of symmetry of the formulation in the presence of explicit antisymmetrization operator $\mathcal{A}_1$. However, this variational property can be used to formulate a basis-set calculational scheme with the following trial functions [14]

$$|\psi_{\mathbf{k}}^1\rangle = \sum_{n=1}^N a_n |f_n\rangle, \quad \langle \psi_{\mathbf{k}}^1| = \sum_{m=1}^N b_m \langle f_m|. \quad (6)$$

Substituting (6) into (5) and using this variational property w.r.t. $|\psi_{\mathbf{k}}^1\rangle$ we obtain [14]

$$\langle \psi_{\mathbf{k}}^1| = \sum_{m=1}^N \langle \phi_{\mathbf{k}}^1 | \mathcal{A}_1 V_1 | f_n \rangle D_{nm} \langle f_m| \quad (7)$$

$$(D^{-1})_{mn} = \langle f_m | \mathcal{A}_1 V_1 - [\mathcal{A}_1 V_1 + (1 - \mathcal{A}_1)(E - H_1)] G_1 \mathcal{A}_1 V_1 | f_n \rangle. \quad (8)$$

Using the variational form (7) and definition $\langle \phi_{\mathbf{k}}^1 | T^{\mathcal{A}} | \phi_{\mathbf{k}}^1 \rangle = \langle \psi_{\mathbf{k}}^1 | \mathcal{A}_1 V_1 | \phi_{\mathbf{k}}^1 \rangle$ we obtain the following basis-set calculational scheme for the transition matrix

$$\langle \phi_{\mathbf{k}}^1 | T^{\mathcal{A}} | \phi_{\mathbf{k}}^1 \rangle = \sum_{m,n=1}^N \langle \phi_{\mathbf{k}}^1 | \mathcal{A}_1 V_1 | f_n \rangle D_{nm} \langle f_m | \mathcal{A}_1 V_1 | \phi_{\mathbf{k}}^1 \rangle. \quad (9)$$

(8) and (9) are also valid for the K matrix and in partial-wave form where the momentum-space integration over the Green's function G_1 should be performed with the principal-value prescription.

In the calculation, the basis functions are taken in the following form

$$f_m(\mathbf{r}_2, \rho_1, \mathbf{s}_1) = \varphi(r_2) \eta(\rho_1) e^{-\delta_m r_2 - \alpha_m \rho_1 - \beta_m s_1 - \gamma_m (\rho_2 + r_{12}) - \mu_m (x + r_1)} \frac{\sin(k s_1)}{k s_1} \quad (10)$$

where $\varphi(r) = \exp(-r)/\sqrt{\pi}$ and $\eta(\rho) = \exp(-0.5\rho)/\sqrt{8\pi}$ represent the H(1s) and Ps(1s) wave functions, respectively. For elastic scattering the direct Born amplitude is zero and the exchange correlation dominates scattering. To be consistent with this, the direct terms in the form factors $\langle f_m | \mathcal{A}_1 V_1 | \phi_{\mathbf{k}}^1 \rangle$ and $\langle \phi_{\mathbf{k}}^1 | \mathcal{A}_1 V_1 | f_n \rangle$ are zero with the above choice of correlations in the basis functions via γ_m and μ_m . This property follows as the above function is invariant w.r.t. the interchange of $\mathbf{x}$ and $\mathbf{r}_1$ whereas the remaining part of the integrand in the direct terms changes sign under this transformation. A proper choice of the correlation parameters γ_m and μ_m is crucial for obtaining good convergence.

In the following we specialize to the K-matrix formulation in S wave at zero energy, when $\sin(k s_1)/(k s_1) = 1$ in (10). The useful matrix elements of the present approach are explicitly written as [14]

$$\langle \phi_p^1 | \mathcal{A}_1 V_1 | f_n \rangle = \pm \frac{1}{2\pi} \int \varphi(\mathbf{r}_1) \eta(\rho_2) \frac{\sin p s_2}{p s_2} [V_1] f_n(\mathbf{r}_2, \rho_1, \mathbf{s}_1) d\mathbf{r}_2 d\rho_1 d\mathbf{s}_1 \quad (11)$$

$$\langle f_m | \mathcal{A}_1 V_1 | \phi_p^1 \rangle = \pm \frac{1}{2\pi} \int f_m(\mathbf{r}_1, \rho_2, \mathbf{s}_2) [V_1] \varphi(\mathbf{r}_2) \eta(\rho_1) \frac{\sin p s_1}{p s_1} d\mathbf{r}_2 d\rho_1 d\mathbf{s}_1 \quad (12)$$

$$\langle f_m | \mathcal{A}_1 V_1 | f_n \rangle = \pm \frac{1}{4\pi} \int f_m(\mathbf{r}_1, \rho_2, \mathbf{s}_2) [V_1] f_n(\mathbf{r}_2, \rho_1, \mathbf{s}_1) d\mathbf{r}_2 d\rho_1 d\mathbf{s}_1 \quad (13)$$

$$\langle f_m | M_1 G_1 \mathcal{A}_1 V_1 | f_n \rangle \approx -\frac{2}{\pi} \int_0^\infty dp \langle f_m | \mathcal{A}_1 V_1 | \phi_p^1 \rangle \langle \phi_p^1 | \mathcal{A}_1 V_1 | f_n \rangle \quad (14)$$

where the so called off-shell term $(1 - \mathcal{A}_1)(E - H_1)$ has been neglected for numerical simplification in this calculation. This term is expected to contribute to refinement over the present calculation. In this convention the on-shell K-matrix element at zero energy is the scattering length: $a = \langle \phi_0^1 | K^{\mathcal{A}} | \phi_0^1 \rangle$.

All the matrix elements above can be evaluated by a method presented in [15]. We describe it in the following for $\langle \phi_p^1 | \mathcal{A}_1 V_1 | f_n \rangle$ of (11). By a transformation of variables from $(\mathbf{r}_2, \rho_1, \mathbf{s}_1)$ to $(\mathbf{s}_1, \mathbf{s}_2, \mathbf{x})$ with Jacobian 2^6 and separating the radial and angular integrations, the form factor (11) is given by

$$\begin{aligned} \langle \phi_p^1 | \mathcal{A}_1 V_1 | f_n \rangle = & \pm \frac{2^6}{16\pi^3} \int_0^\infty s_2^2 ds_2 \frac{\sin(ps_2)}{ps_2} \int_0^\infty s_1^2 ds_1 e^{-\beta_n s_1} \int_0^\infty x^2 dx e^{-\mu_n x} \\ & \times \int e^{-(ar_1+b\rho_1/2)} e^{-(cr_2+d\rho_2/2)} e^{-\gamma_n r_{12}} [V_1] d\hat{s}_1 d\hat{s}_2 d\hat{x} \end{aligned} \quad (15)$$

where $a = 1 + \mu_n$, $b = 2\alpha_n + 1$, $c = 1 + \delta_n$ and $d = 2\gamma_n + 1$. Recalling that $\mathbf{r}_j = 2\mathbf{s}_j - \mathbf{x}$, $\mathbf{r}_{12} = 2(\mathbf{s}_1 - \mathbf{s}_2)$, $\rho_j = 2(\mathbf{x} - \mathbf{s}_j)$, $j = 1, 2$, we employ the following expansions of the exponentials in (15)

$$e^{-a|2\mathbf{s}-\mathbf{x}|-b|\mathbf{x}-\mathbf{s}|} = \frac{4\pi}{sx} \sum_{lm} G_l^{(a,b)}(s, x) Y_{lm}^*(\hat{s}) Y_{lm}(\hat{x}) \quad (16)$$

$$\frac{e^{-a|2\mathbf{s}-\mathbf{x}|-b|\mathbf{x}-\mathbf{s}|}}{|2\mathbf{s}-\mathbf{x}|} = \frac{4\pi}{sx} \sum_{lm} J_l^{(a,b)}(s, x) Y_{lm}^*(\hat{s}) Y_{lm}(\hat{x}) \quad (17)$$

$$\frac{e^{-a|2\mathbf{s}-\mathbf{x}|-b|\mathbf{x}-\mathbf{s}|}}{|\mathbf{s}-\mathbf{x}|} = \frac{4\pi}{sx} \sum_{lm} K_l^{(a,b)}(s, x) Y_{lm}^*(\hat{s}) Y_{lm}(\hat{x}) \quad (18)$$

$$\frac{e^{-a|\mathbf{s}_1-\mathbf{s}_2|}}{|\mathbf{s}_1-\mathbf{s}_2|} = \frac{4\pi}{s_1 s_2} \sum_{lm} A_l^{(a)}(s_1, s_2) Y_{lm}^*(\hat{s}_1) Y_{lm}(\hat{s}_2) \quad (19)$$

$$e^{-a|\mathbf{s}_1-\mathbf{s}_2|} = \frac{4\pi}{s_1 s_2} \sum_{lm} B_l^{(a)}(s_1, s_2) Y_{lm}^*(\hat{s}_1) Y_{lm}(\hat{s}_2) \quad (20)$$

where the Y_{lm} 's are the usual spherical harmonics. Using (16) – (20) in (15) we get

$$\begin{aligned} \langle \phi_p^1 | \mathcal{A}_1 V_1 | f_n \rangle = & \pm 2^8 \int_0^\infty e^{-\beta_n s_1} ds_1 \int_0^\infty ds_2 \frac{\sin(ps_2)}{ps_2} \int_0^\infty dx e^{-\mu_n x} \sum_{l=0}^L (2l+1) \\ & \times \left[\frac{1}{x} G_l^{(a,b)}(s_1, x) G_l^{(c,d)}(s_2, x) B_l^{(2\gamma_n)}(s_1, s_2) - J_l^{(a,b)}(s_1, x) \right. \\ & \times G_l^{(c,d)}(s_2, x) B_l^{(2\gamma_n)}(s_1, s_2) + \frac{1}{2} G_l^{(a,b)}(s_1, x) G_l^{(c,d)}(s_2, x) \\ & \left. \times A_l^{(2\gamma_n)}(s_1, s_2) - \frac{1}{2} G_l^{(a,b)}(s_1, x) K_l^{(c,d)}(s_2, x) B_l^{(2\gamma_n)}(s_1, s_2) \right]. \end{aligned} \quad (21)$$

where the l -sum is truncated to $l = L$. This evaluation avoids complicated angular integrations involving $\mathbf{s}_1$, $\mathbf{s}_2$ and $\mathbf{x}$. These integrals take a simple form requiring straightforward numerical computation of certain radial integrals only, which must, however, be carried out carefully. The functions G_l , J_l , K_l etc. are easily calculated using (16) – (20):

$$G_l^{(a,b)}(s, x) = \frac{sx}{2} \int_{-1}^1 dt P_l(t) e^{-a|2\mathbf{s}-\mathbf{x}|-b|\mathbf{x}-\mathbf{s}|} \quad (22)$$

where $P_l(t)$ is the usual Legendre polynomial and t is the cosine of the angle between $\mathbf{s}$ and $\mathbf{x}$. The integrals (12) and (13) can be evaluated similarly. For example

$$\begin{aligned} \langle f_m | \mathcal{A}_1 V_1 | \phi_p^1 \rangle &= \pm 2^8 \int_0^\infty e^{-\beta_m s_2} ds_2 \int_0^\infty ds_1 \frac{\sin(ps_1)}{ps_1} \int_0^\infty dx e^{-\mu_m x} \sum_{l=0}^L (2l+1) \\ &\times \left[\frac{1}{x} G_l^{(c,d)}(s_1, x) G_l^{(a,b)}(s_2, x) B_l^{(2\gamma_m)}(s_1, s_2) - J_l^{(c,d)}(s_1, x) \right. \\ &\times G_l^{(a,b)}(s_2, x) B_l^{(2\gamma_m)}(s_1, s_2) + \frac{1}{2} G_l^{(c,d)}(s_1, x) G_l^{(a,b)}(s_2, x) \\ &\times A_l^{(2\gamma_m)}(s_1, s_2) - \left. \frac{1}{2} G_l^{(c,d)}(s_1, x) K_l^{(a,b)}(s_2, x) B_l^{(2\gamma_m)}(s_1, s_2) \right] \quad (23) \end{aligned}$$

$$\begin{aligned} \langle f_m | \mathcal{A}_1 V_1 | f_n \rangle &= \pm 2^7 \int_0^\infty e^{-\beta_n s_1} ds_1 \int_0^\infty ds_2 e^{-\beta_m s_2} \int_0^\infty dx e^{-(\mu_n + \mu_m)x} \sum_{l=0}^L (2l+1) \\ &\times \left[\frac{1}{x} G_l^{(e,f)}(s_1, x) G_l^{(g,h)}(s_2, x) B_l^{(2\gamma_{mn})}(s_1, s_2) - J_l^{(e,f)}(s_1, x) \right. \\ &\times G_l^{(g,h)}(s_2, x) B_l^{(2\gamma_{mn})}(s_1, s_2) + \frac{1}{2} G_l^{(e,f)}(s_1, x) G_l^{(g,h)}(s_2, x) \\ &\times A_l^{(2\gamma_{mn})}(s_1, s_2) - \left. \frac{1}{2} G_l^{(e,f)}(s_1, x) K_l^{(g,h)}(s_2, x) B_l^{(2\gamma_{mn})}(s_1, s_2) \right] \quad (24) \end{aligned}$$

where $e = 1 + \delta_m + \mu_n$, $f = 2\alpha_n + 2\gamma_m + 1$, $g = 1 + \delta_n + \mu_m$, $h = 2\alpha_m + 2\gamma_n + 1$ and $\gamma_{mn} = \gamma_m + \gamma_n$.

We tested the convergence of the integrals by varying the number of integration points in the x , s_1 and s_2 integrals in (21), (23) and (24) and the t integral in (22). The evaluation of (21), (23) and (24) essentially involves four-dimensional integration, which is performed with caution. The x integration was relatively easy and 20 Gauss-Legendre quadrature points appropriately distributed between 0 and 16 were enough for convergence. In the evaluation of integrals of type (22) 40 Gauss-Legendre quadrature points were sufficient for adequate convergence. The convergence in the numerical integration over s_1 and s_2 was achieved with 300 Gauss-Legendre quadrature points between 0 and 12. The maximum value of l

in the sum in (21), (23) and (24), L , is taken to be 6 which is sufficient for obtaining the convergence with the partial-wave expansions (16) – (20).

Table 1: Singlet (a_s) and triplet (a_t) Ps-H scattering lengths for different L and N .

N	$L = 0$		$L = 2$		$L = 4$		$L = 6$	
	a_t	a_s	a_t	a_s	a_t	a_s	a_t	a_s
6	4.00	3.61	3.66	4.00	3.07	3.75	2.92	3.74
7	3.98	4.11	3.25	3.88	2.99	4.06	2.85	4.00
8	3.93	4.12	3.35	3.96	2.66	3.72	2.54	3.73
9	3.97	4.15	3.42	3.83	2.65	3.92	2.55	4.06
10	3.97	4.22	3.42	3.92	2.57	4.42	2.50	3.45
11	4.01	3.82	3.44	3.91	3.55	3.75	2.67	3.80
12					2.54	3.72	2.46	3.73
13					2.48	3.47	2.46	3.49

In a numerical calculation a judicious choice of the parameters in (10) is needed for rapid convergence. As the variational method does not provide a bound on the result, the method could converge to a wrong scattering length if an inappropriate (incomplete) basis set is chosen. After some experimentation we find that to obtain proper convergence the parameters δ_n and α_n should be taken to have both positive and negative values, γ_n and μ_n should include values close to unity and β_n should have progressively increasing values till about 1.5. The results reported in this work are obtained with the following parameters for the functions $f_n, n = 1, \dots, 13$: $\{\delta_n, \alpha_n, \beta_n, \gamma_n, \mu_n\} \equiv \{-0.5, -0.25, 0.3, 0.01, 0.02\}, \{-0.5, -0.25, 0.5, 0.04, 0.02\}, \{-0.5, -0.25, 0.7, 0.03, 0.06\}, \{-0.2, -0.1, 0.6, 0.2, 0.2\}, \{-0.1, 0.1, 0.8, 0.25, 0.25\}, \{0.2, -0.2, 0.6, 0.35, 0.35\}, \{-0.1, -0.1, 0.7, 0.4, 0.4\}, \{0.15, 0.2, 0.8, 0.5, 0.5\}, \{0.12, -0.12, 1, 0.7, 0.7\}, \{0.2, 0.2, 1.2, 0.9, 0.9\}, \{0.1, 0.2, 1.3, 1, 0.7\}, \{0.2, 0.1, 1.4, 0.7, 1\}, \{0.3, 0.15, 1.5, 1, 1\}$

In table 1 we show the convergence pattern of the present calculation w.r.t. the number of partial waves L and basis functions N used in the calculation. The convergence is smooth with increasing L . However, as the present calculation does not produce a bound on the

result, the convergence is not monotonic with increasing N . The lack of an upper bound in this calculation is clearly revealed in table 1 where the results do not decrease monotonically as the number of terms in the trial wave function is increased. The unbounded nature of the results is consistent with the noted oscillation of the scattering lengths as N increases. The oscillation is larger in the singlet state where it is more difficult to obtain convergence. Although, the final result for the largest N and L is supposed to be the most accurate, it is not known whether this result is larger or smaller than the exact one. Also, an estimate of error of this result is not known. It is difficult to provide a quantitative measure of convergence. However, from the noted fluctuation of the results for large N for $L = 6$ we believe the error in the triplet scattering length to be less than 0.10 and in the singlet scattering length to be less than 0.20. The final results of the present calculation are those for $N = 13$ and $L = 6$ with the above estimate of error: $a_s = 3.49 \pm 0.20$ au and $a_t = 2.46 \pm 0.10$ au.

The maximum number of functions ($N = 13$) used in this calculation is also pretty small, compared to those used in different Kohn-type variational calculations for electron-hydrogen ($N = 56$) [16] and positron-hydrogen ($N \leq 286$) [17] scattering. Because of the explicit appearance of the Green's function, the present basis-set approach is similar to the Schwinger variational method. Using the Schwinger method, convergent results for electron-hydrogen [18] and positron-hydrogen [19] scattering have been obtained with a relatively small basis set ($N \sim 10$). These suggest a more rapid convergence in these problems with a Schwinger-type method.

Now we compare the present results with those of other calculations. While a static-exchange calculation by Hara and Fraser [10] yielded $a_s = 7.28$ au and $a_t = 2.48$ au a model potential calculation by Drachman and Houston [11] produced $a_s = 5.33$ au and $a_t = 2.36$ au. A 22-state R-matrix calculation by Campbell et al. [3] and a six-state CC calculation by Sinha et al. [4] yielded $a_s = 5.20$ au, $a_t = 2.45$ au and $a_s = 5.90$ au, $a_t = 2.32$ au, respectively. These triplet scattering lengths are in qualitative agreement with the present result: $a_t = 2.46 \pm 0.10$. However, the singlet scattering lengths of these calculations have not yet converged. The disagreement in the singlet channel shows that it is more difficult to get the converged result in the attractive singlet channel than in the repulsive

triplet channel. This is consistent with the common wisdom that a scattering model at low energies is more sensitive to the detail of an effective attractive interaction than to that of a repulsive interaction. The nonconvergence of these results for the singlet scattering length was conjectured before [5]. Using a correlation between the S-wave scattering length and binding energy for the Ps-H system, the value $a_s = 3.5$ au was predicted in that study [5] in excellent agreement with the present result: $a_s = 3.49 \pm 0.20$.

To summarize, we have formulated a convergent basis-set calculational scheme for S-wave Ps-H elastic scattering below the lowest inelastic threshold using a variational expression for the transition matrix. We illustrate the method numerically by calculating the singlet and triplet scattering lengths: $a_s = 3.49 \pm 0.20$ au and $a_t = 2.46 \pm 0.10$ au.

The work is supported in part by the Conselho Nacional de Desenvolvimento - Científico e Tecnológico, Fundação de Amparo à Pesquisa do Estado de São Paulo, and Financiadora de Estudos e Projetos of Brazil.

REFERENCES

- [1] Garner A J, Laricchia G and Özen A 1996 *J. Phys. B: At. Mol. Opt. Phys.* **29** 5961
Garner A J, Özen A and Laricchia G 2000 *J. Phys. B: At. Mol. Opt. Phys.* **33** 1149
Skalsey M, Engbrecht J J, Bithell R K, Vallery R S and Gidley D W 1998 *Phys. Rev. Lett.* **80** 3727
Nagashima Y, Hyodo T, Fujiwara K and Ichimura A 1998 *J. Phys. B: At. Mol. Opt. Phys.* **31** 329
- [2] Andersen H H, Armour E A G, Humberston J W and Laricchia G 1998 *Nucl. Instrum. & Methods Phys. Res. B* **143** U10
Biswas P K 2000 *Nucl. Instrum. & Methods Phys. Res. B* **171** 135
- [3] Campbell C P, McAlinden M T, MacDonald F G R S and Walters H R J 1998 *Phys. Rev. Lett.* **80** 5097
- [4] Sinha P K, Basu A and Ghosh A S 2000 *J. Phys. B: At. Mol. Opt. Phys.* **33** 2579
- [5] Adhikari S K and Biswas P K 1999 *Phys. Rev. A* **59** 2058
Adhikari S K 2001 *Phys. Rev. A* **63** in press
- [6] Biswas P K and Adhikari S K 1999 *Phys. Rev. A* **59** 363
Adhikari S K 2000 *Phys. Rev. A* **62** 062708
- [7] Biswas P K and Adhikari S K 2000 *Chem. Phys. Lett.* **317** 129
- [8] Biswas P K and Adhikari S K 2000 *J. Phys. B: At. Mol. Opt. Phys.* **33** 1575
- [9] Barker M I and Bransden B H 1968 *J. Phys. B: At. Mol. Opt. Phys.* **1** 1109
- [10] Hara S and Fraser P A 1975 *J. Phys. B: At. Mol. Opt. Phys.* **8** L472
- [11] Drachman R J and Houston S K 1975 *Phys. Rev. A* **12** 885
Drachman R J and Houston S K 1976 *Phys. Rev. A* **14** 894
- [12] Frolov A M and Smith Jr. V H 1997 *Phys. Rev. A* **55** 2662
Yan Z C and Ho Y K 1999 *Phys. Rev. A* **59** 2697

- Jiang N and Schrader D M 1998 *J. Chem. Phys.* **109** 9430
- [13] Adhikari S K 1982 *Phys. Rev. C* **25** 118
- [14] Adhikari S K and Sloan I H 1975 *Phys. Rev. C* **11** 1133
- Adhikari S K and Sloan I H 1975 *Nucl. Phys.* **A241** 429
- Adhikari S K 1974 *Phys. Rev. C* **10** 1623
- Adhikari S K 1998 *Variational Principles and the Numerical Solution of Scattering Problems* (New York: John-Wiley)
- [15] Adhikari S K and Mandal P 2000 *J. Phys. B: At. Mol. Opt. Phys.* **33** L761
- [16] Shimamura I 1971 *J. Phys. Soc. Japan* **30** 1702
- [17] Humberston J W, VanReeth P, Watts M S T and Meyerhof W E 1997 *J. Phys. B: At. Mol. Opt. Phys.* **30** 2477
- Armour E A G and Humberston J W 1991 *Phys. Rep.* **204** 1734
- [18] Takatsuka K and McKoy V 1984 *Phys. Rev. A* **30** 1734
- [19] Kar S and Mandal P 2000 *J. Phys. B: At. Mol. Opt. Phys.* **33** 2379
- Kar S and Mandal P 1999 *J. Phys. B: At. Mol. Opt. Phys.* **32** 2297
- Kar S and Mandal P 1999 *Phys. Rev. A* **59** 1913

Figure Caption:

1. Different position vectors for the Ps-H system w.r.t the proton (p) at the origin in arrangement 1 with electron 1 forming the Ps.




Figure 1